

CONTENTS


	Pages
ABSTRACT (Thai)	i
ABSTRACT (English)	iii
ACKNOWLEDGEMENTS	v
CONTENTS	vi
LIST OF TABLES	viii
LIST OF FIGURES	ix
CHAPTERS	
CHAPTER I	
INTRODUCTION	
- Cholangiocarcinoma	1
- Apoptosis	3
- Tumor Necrosis Factor alpha (TNF-alpha)	5
CHAPTER II	
LITERATURE REVIEW	10
CHAPTER III	
OBJECTIVES	13
CHAPTER IV	
MATERIALS AND METHODS	
- Cell culture: thawing and recovering cell line	14
- Trypsinization and subculture method	14
- Determination of TNFR mRNA expression by RT-PCR	15

CONTENTS

	Pages
- Determination of apoptosis cells by DAPI staining	17
- Determination of cytotoxicity by MTT assay	18
- Determination of poly (ADP ribose) polymerase (PARP) by Western Blot Assay	19
CHAPTER V RESULTS	
- Tumor Necrosis Factor Receptor (TNFR) expression	23
- DAPI staining assay	25
- MTT assay	44
- Western Blot assay	47
CHAPTER VI DISCUSSION	49
CHAPTER VII CONCLUSION	54
REFERENCES	55
APPENDIX	
- Chemical substances and equipment	61
- Reagents	63
- Statistical analysis	68
BIOGRAPHY	71

LIST OF TABLES

	Pages
Table 1 Cytotoxicity of KKU-100 performed by using MTT assay represent in percentage.	45
Table 2 Cytotoxicity of KKU-213 perform by using MTT assay represent in percentage.	46
Table 3 Multiple Comparisons of MTT assay of KKU-100 cell line.	68
Table 4 Multiple Comparisons of MTT assay of KKU-213 cell line.	69



LIST OF FIGURES

	Pages
Figure 1 Age-standardized incidence rates in European and Asian populations by gender.	2
Figure 2 Schematic overview of the major apoptosis pathways and their inhibitors.	4
Figure 3 Signal transduction pathway initiated by trimeric TNF-alpha binding to its receptor, TNFR to initiate receptor clustering and signal transduction.	9
Figure 4 Expression of TNFRI (upper) and TNFRII (lower) of the KKU-213 and KKU-100 cell lines was determined by RT-PCR.	24
Figure 5 DAPI staining analysis (First experiment): KKU-100 cells were seeded onto coverslips for 24 h and absent of 160 ng/ml TNF-alpha (control condition).	26
Figure 5.1 DAPI staining analysis (second experiment): KKU-100 cells stained with DAPI in control condition (absent of TNF-alpha).	27
Figure 5.2 DAPI staining analysis (third experiment): KKU-100 cells stained with DAPI in control condition (absent of TNF-alpha).	28
Figure 6 DAPI staining analysis (first experiment): KKU-100 cells stained with DAPI in TNF-alpha condition (present with 160 ng/ml TNF-alpha for 24 h).	29
Figure 6.1 DAPI staining analysis (second experiment): KKU-100 cells stained with DAPI in TNF-alpha condition (present with 160 ng/ml TNF-alpha for 24 h).	30

LIST OF FIGURES

	Pages
Figure 6.2 DAPI staining analysis (first experiment): KKU-100 cells stained with DAPI in TNF-alpha condition (present with 160 ng/ml TNF-alpha for 24 h).	31
Figure 7 DAPI staining analysis (first experiment): KKU-100 cells stained with DAPI in positive control condition (present with 15 mM Wortmanin for 24 h).	32
Figure 7.1 DAPI staining analysis (second experiment): KKU-100 cells stained with DAPI in positive control condition (present with 15 mM Wortmanin).	33
Figure 7.2 DAPI staining analysis (third experiment): KKU-100 cells stained with DAPI in positive control condition (present with 15 mM Wortmanin).	34
Figure 8 DAPI staining analysis (first experiment): KKU-213 cells stained with DAPI in control condition (absent of TNF-alpha).	35
Figure 8.1 DAPI staining analysis (second experiment): KKU-213 cells stained with DAPI in control condition (absent of TNF-alpha).	36
Figure 8.2 DAPI staining analysis (third experiment): KKU-213 cells stained with DAPI in control condition (absent of TNF-alpha).	37
Figure 9 DAPI staining analysis (first experiment): KKU-213 cells stained with DAPI in TNF-alpha condition (present of 160 ng/ml TNF-alpha).	38

LIST OF FIGURES

	Pages
Figure 9.1 DAPI staining analysis (second experiment): KKU-213 cells stained with DAPI in TNF-alpha condition (present of 160 ng/ml TNF-alpha).	39
Figure 9.2 DAPI staining analysis (third experiment): KKU-213 cells stained with DAPI in TNF-alpha condition (present of 160 ng/ml TNF-alpha).	40
Figure 10 DAPI staining analysis (first experiment): KKU-213 cells stained with DAPI in positive control condition (present with 15 mM Wortmanin).	41
Figure 10.1 DAPI staining analysis (second experiment): KKU-213 cells stained with DAPI in positive control condition (present with 15 mM Wortmanin).	42
Figure 10.2 DAPI staining analysis (third experiment): KKU-213 cells stained with DAPI in positive control condition (present with 15 mM Wortmanin).	43
Figure 11 Cytotoxicity of KKU100 after 160 ng/ml TNF-alpha treatment, compared with control (untreated) and positive control treated with 15 mM Wortmanin.	45
Figure 12 Cytotoxicity of KKU-213 after 160 ng/ml TNF-alpha treatment compared with control (untreated) and positive control treated with 15 mM Wortmanin.	46
Figure 13 Western blot analysis of PARP fragments of KKU-100 (A) and KKU-213 (B) cell lines.	48