

บรรณานุกรม

- Ajo, D. and Casarin, M. et al (1981) Conformational flexibility of the dehydrophenylalanine derivatives: Molecular and electronic structure of (*Z*)-N-acetyl dehydrophenylalanine. *Tetrahedron*, 37 : 3507-3512.
- Braun, S. and Raymond, W. E. et al (1984) Synthetic tyrosine polymers as substrates and inhibitors of tyrosine-specific protein kinases. *J. Biol. Chem.*, 259 : 2051-2054.
- Brunton, V.G. and Workman, P. (1993) Cell signaling targets for antitumour drug development. *Cancer Chemother. Pharmacol.*, 32 : 1-19.
- Burke, R.T. (1992) Protein-tyrosine kinase inhibitors. *Drug of The Future*, 17 : 119-131.
- Burke, T. R.Jr., and Lim, B. et al (1993) Bicyclic compounds as ring-constrained inhibitors of protein-tyrosine kinase^{PS6001}. *J. Med. Chem.*, 36 : 425-431.
- Burke, T. R.Jr and Li, Z. et.al (1991) Phosphonate-containing inhibitors of tyrosine-specific protein kinase. *J. Med. Chem.*, 34 : 1577-1581.
- Chang, C. and Geahlen, R. L.(1992) Protein-tyrosine kinase inhibition; mechanism-based discovery of antitumor agents. *J. of Natural Products* 1, 55 : 1529-1560.
- Gazit, A., Yaish, P. and Levitzki, A. Tyrphostins I: synthesis and biological activity of protein tyrosine kinase inhibitors. *J. Med. Chem.* 3 : 2344-2352.
- Kenyon , G. I. and Garcia, G.A.(1987) Design of kinase inhibitors. *Med. Res. Rev.* 7 : 389-416.
- Kruse, C. H. and Holden, K. G. et al. (1988) Synthesis and evaluation of multisubstrate inhibitors of an oncogene-encoded tyrosine specific protein kinase. 2 . *J. Med. Chem.*, 31 : 1768-1772.
- Latif, N. and Mishriky, N.; et al (1982) Reaction of biologically active β -nitrostyrenes with *o*-phenylenediamine:a new route to the synthesis of substituted benzimidazoles. *Indian J. Chem. Sect. B*, 21 : 872-874.
- Reidel, H. and Dull, T. J. et al (1986) A chimeric receptor allows insulin to stimulate tyrosine kinase activity of epidermal growth factor receptor. *Nature* , 324 : 68-70.
- Shechter, H., Ley, D. E. and Roberson, E. B. Jr. (1956) Nitroalkanes from conjugated nitroalkenes by reduction with complex hydrides. *J. Am. Chem. Soc.*, 78 : 4984-4991.
- Shiraishi, T. and Owada, M. K. et al. (1989) Specific inhibitors of tyrosine-specific protein kinases, properties of 4-Hydroxycinnamamide derivatives *in vitro*. *Cancer Res.* 49 : 2374-2378.
- Trexler, P M. and Wacker, O. et al (1991) Sulfonylbenzoyl-nitrostyrene: potential bisubstrate type inhibitors of the EGF receptor tyrosine protein kinase. *J. Med. Chem.* 34 : 2328-2337.

Wollenberg, R. H. and Miller, S. J. (1978) Nitroalkanes synthesis. A convenient method for aldehyde reductive nitromethylation. *Tet. Lett.*, 35 : 3219-3222.

Workman, P., Bruton, V.G. and Robins, D.J. (1992) Tyrosine kinase inhibitors. *Seminars in Cancer Biology*, 3: 369-381.

Yaish, P. and Gazit, A. et al (1988) Blocking of EGF-dependent cell proliferation by EGF receptor kinase inhibitors. *Science*, 242 : 933-935.



ประวัติย่อผู้วิจัย

ชื่อ-นามสกุล

ดร.อนงค์ ตีระวนิชพงศ์

ประวัติการศึกษา

เก็ชชานาสตร์บัณฑิต (ก.บ.), มหาวิทยาลัยเชียงใหม่
Ph.D.(Medicinal Chemistry), Duquesne University, Pittsburgh, PA, USA.

สถานที่ติดต่อ

สาขาวิชาเภสัชเวชและเภสัชเคมี คณะเภสัชศาสตร์
มหาวิทยาลัยหัวเฉียวเฉลิมพระเกียรติ
โทรศัพท์ 312-6300 ต่อ 1494, 1171
e-mail : anong@hcu.ac.th

